

## ***When Immunostains Can Get You in Trouble: Gynecologic Pathology p16: Panacea or Pandora's Box?***

### REFERENCES

1. Darragh TM, Colgan TJ, Cox JT, et al. The Lower Anogenital Squamous Terminology Standardization Project for HPV-Associated Lesions: background and consensus recommendations from the College of American Pathologists and the American Society for Colposcopy and Cervical Pathology. *Arch Pathol Lab Med.* 2012;136:1266-1297.
2. Raju RR GJ, Hart WR. Pagetoid squamous cell carcinoma in situ (pagetoid Bowen's disease) of the external genitalia. *Int J Gynecol Pathol.* 2003;22:127-135.
3. Longacre TA KC, Welton ML. Diagnostic problems in anal pathology. *Adv Anat Pathol.* 2008;15:263-278.
4. Bala R, Pinsky BA, Beck AH, et al. p16 is superior to ProEx C in identifying high-grade squamous intraepithelial lesions (HSIL) of the anal canal. *Am J Surg Pathol.* 2013;37:659-668.
5. Darragh TM CT, Thomas Cox J, Heller DS, Henry MR, Luff RD, McCalmont T, Nayar R, Palefsky JM, Stoler MH, Wilkinson EJ, Zaino RJ, Wilbur DC, Groups MotLPW. The Lower Anogenital Squamous Terminology Standardization project for HPV-associated lesions: background and consensus recommendations from the Col... - PubMed - NCBI. *Int J Gynecol Pathol.* 2013;32:76-115.
6. Keating JT, Cviko A, Riethdorf S, et al. Ki-67, cyclin E, and p16INK4 are complimentary surrogate biomarkers for human papilloma virus-related cervical neoplasia. *Am J Surg Pathol.* 2001;25:884-891.
7. Iaconis L, Hyjek E, Ellenson LH, et al. p16 and Ki-67 immunostaining in atypical immature squamous metaplasia of the uterine cervix: correlation with human papillomavirus detection. *Arch Pathol Lab Med.* 2007;131:1343-1349.
8. Qiao X, Bhuiya TA, Spitzer M. Differentiating high-grade cervical intraepithelial lesion from atrophy in postmenopausal women using Ki-67, cyclin E, and p16 immunohistochemical analysis. *J Low Genit Tract Dis.* 2005;9:100-107.
9. Hoevenaars BM, van der Avoort IA, de Wilde PC, et al. A panel of p16(INK4A), MIB1 and p53 proteins can distinguish between the 2 pathways leading to vulvar squamous cell carcinoma. *Int J Cancer.* 2008;123:2767-2773.
10. Kong CS, Balzer BL, Troxell ML, et al. p16INK4A immunohistochemistry is superior to HPV in situ hybridization for the detection of high-risk HPV in atypical squamous metaplasia. *Am J Surg Pathol.* 2007;31:33-43.
11. Pocrnich CE, Ramalingam P, Euscher ED, et al. Neuroendocrine carcinoma of the endometrium: a clinicopathologic study of 25 cases *Am J Surg Pathol.* 2016;40:577-86
12. Mills AM, Ly A, Balzer BL, et al. Cell cycle regulatory markers in uterine atypical leiomyoma and leiomyosarcoma: immunohistochemical study of 60 cases with clinical follow-up. *Am J Surg Pathol,* 2013;37:634-42.
13. Kong CS, Beck AH, Longacre TA. A panel of three markers including p16, ProEx C, or HPV ISH is optimal for distinguishing between primary endometrial and endocervical adenocarcinomas *Am J Surg Pathol.* 2010;34:915-26.

14. Atkins KA, Arronte N, Darus CJ, et al. The Use of p16 in enhancing the histologic classification of uterine smooth muscle tumors. *Am J Surg Pathol*. 2008;32:98-102.
15. O'Neill CJ, McBride HA, Connolly LE, et al. Uterine leiomyosarcomas are characterized by high p16, p53 and MIB1 expression in comparison with usual leiomyomas, leiomyoma variants and smooth muscle tumours of uncertain malignant potential. *Histopathology*. 2007;50:851-858.
16. Bodner-Adler B, Bodner K, Czerwenka K, et al. Expression of p16 protein in patients with uterine smooth muscle tumors: an immunohistochemical analysis. *Gynecol Oncol*. 2005;96:62-66.
17. Skubitz KM, Skubitz AP. Differential gene expression in leiomyosarcoma. *Cancer*. 2003;98:1029-1038.
18. Unver NU, Acikalin MF, Oner U, et al. Differential expression of P16 and P21 in benign and malignant uterine smooth muscle tumors. *Arch Gynecol Obstet*. 2011;284:483-490.
19. Park KJ, Kiyokawa T, Soslow RA, et al. Unusual endocervical adenocarcinomas: an immunohistochemical analysis with molecular detection of human papillomavirus. *Am J Surg Pathol*. 2011;35:633-46.
20. Houghton O, Jamison J, Wilson R, et al. p16 Immunoreactivity in unusual types of cervical adenocarcinoma does not reflect human papillomavirus infection. *Histopathology*. 2010;57:342-50.
21. Ordi J, Alejo M, Fuste V, et al. HPV-negative vulvar intraepithelial neoplasia (VIN) with basaloid histologic pattern: an unrecognized variant of simplex (differentiated) VIN. *Am J Surg Pathol*. 2009;33:1659-1665
22. van Seters M, ten Kate FJW, M van Beurden M, et al. In the absence of (early) invasive carcinoma, vulvar intraepithelial neoplasia associated with lichen sclerosus is mainly of undifferentiated type: new insights in histology and aetiology. *J Clin Pathol* 2007;60:504-509
23. Brainard JA, Hart WR. Proliferative epidermal lesions associated with anogenital Paget's disease. *Am J Surg Pathol*. 2000;24:543-52.
24. Int J Gynecol Pathol. 2012;31:206-10.
25. McCluggage WG, McBride HA. Papillary syncytial metaplasia associated with endometrial breakdown exhibits an immunophenotype that overlaps with uterine serous carcinoma. *Int J Gynecol Pathol*. 2012;31:206-10
26. Yemelyanova A, Ji H, Shih IeM, et al. Utility of p16 expression for distinction of uterine serous carcinomas from endometrial endometrioid and endocervical adenocarcinomas: immunohistochemical analysis of 201 cases. *Am J Surg Pathol* 2009;33:1504-14.
27. Chiesa-Vottero AG, Malpica A, Deavers MT, et al. Immunohistochemical overexpression of p16 and p53 in uterine serous carcinoma and ovarian high-grade serous carcinoma. *Int J Gynecol Pathol* 2007;26:328-33
28. Alexander RE, Hu Y, Kum JB, Montironi R, et al. p16 expression is not associated with human papillomavirus in urinary bladder squamous cell carcinoma. *Mod Pathol* 2012;25:1526-1533.

29. Alexander RE, Williamson SR, Richey J. et al. The expression patterns of p53 and p16 and an analysis of a possible role of HPV in primary adenocarcinoma of the urinary bladder. *PLoS One* 2014; 9:e95724
30. Schwartz LE, Khani F, Bishop JA, et al. Carcinoma of the uterine cervix involving the genitourinary tract: a potential diagnostic dilemma. *Am J Surg Pathol* 2016;40:27-35